

Solution to worldvolume action of D3 brane in pp-wave background

Shesansu Sekhar Pal
e-mail: shesansu@iopb.res.in

*Institute of Physics
Bhubaneswar - 751005, India*

Abstract

We find nonsupersymmetric and supersymmetric solutions of D3 brane configuration in the background of pp wave obtained as a Penrose limit of $AdS_5 \times S^5$.

Recently, it has been shown that the study of maximally supersymmetric IIB supergravity background, pp waves[1], of various geometries gives rise to many interesting ideas in the context of AdS/CFT correspondence. Especially, the study of BMN[2] tells us how to see the duality in a particular sector of the four dimensional SU(N) gauge theory with $\mathcal{N} = 4$ with type IIB superstring theory in pp wave background. The study of duality¹ have been extended to various singular spaces with different supersymmetries in [26]-[35] and have been studied in [31], [32], [33] from holographic point of view. Moreover, string theory in this background is exactly solvable. The quantization of strings in the pp wave background with NS-NS and R-R fluxes is done in [36]-[39] and the quantization of Dp brane is studied in [40]. Solution to branes in the pp wave background have been studied in [41]-[42] and in [43],[44] from supergravity point of view. The authors of [42] have studied various configurations of Dp brane with different embeddings and found solutions of constant embeddings with zero worldvolume flux on the brane.

In this letter we shall derive nonsupersymmetric and supersymmetric solutions of D3 brane whose worldvolume directions are extended along any one of the SO(4) directions and one along positive lightcone direction and others are along one of the SO(4) directions in the pp wave background.

Before we start to present the solution of D3 brane in the pp wave background, let us discuss the geometry briefly. The geometry of $AdS_p \times S^q$ in global coordinate can be described as:

$$ds^2 = R_A^2(d\rho^2 + \sinh^2 \rho d\Omega_{p-2}^2 - \cosh^2 \rho dt^2) + R_S^2(d\theta^2 + \sin^2 \theta d\Omega_{q-2}^2 + \cos^2 \theta d\psi^2), \quad (1)$$

where R_A, R_S are the radius of curvature and radius of AdS_p and S^q respectively. To derive the pp wave, we have to consider a particle moving along the ψ direction and sitting at $\rho = 0$ and $\theta = 0$ and the geometry seen by the particle while moving along this trajectory will give us the desired geometry. To do so, let us define coordinates as:

$$x^+ = \frac{1}{2}(t + \frac{R_S}{R_A}\psi), \quad x^- = \frac{R_A^2}{2}(t - \frac{R_S}{R_A}\psi), \quad x = R_A\rho, \quad y = R_S\theta. \quad (2)$$

Hence, the geometry in this coordinates becomes

$$ds^2 = R_A^2 \left[\frac{dx^2}{R_A^2} + \sinh^2 \left(\frac{x}{R_A} \right) d\Omega_{p-2}^2 - \cosh^2 \left(\frac{x}{R_A} \right) \left\{ (dx^+)^2 + R_A^{-4} (dx^-)^2 + 2R_A^{-2} dx^+ dx^- \right\} \right] \\ + R_S^2 \left[\frac{dy^2}{R_S^2} + \sin^2 \left(\frac{y}{R_S} \right) d\Omega_{q-2}^2 + \cos^2 \left(\frac{y}{R_S} \right) \left\{ \frac{R_A^2 (dx^+)^2}{R_S^2} + \frac{(dx^-)^2}{R_S^2 R_A^2} - \frac{2}{R_S^2} dx^+ dx^- \right\} \right] \quad (3)$$

Let us define a limit

$$R_A \rightarrow \infty, \quad R_S \rightarrow \infty, \quad \text{and} \quad \text{keeping} \quad \frac{R_S}{R_A} = \text{fixed}. \quad (4)$$

¹Also studied in [3], see also refs. [4]-[25] for related recent developments.

In this limit the metric becomes

$$ds^2 = -4dx^+dx^- - \left(x^2 + \frac{R_A^2}{R_S^2}y^2\right)(dx^+)^2 + dx^2 + x^2d\Omega_{p-2}^2 + dy^2 + y^2d\Omega_{q-2}^2, \quad (5)$$

scaling $x^+ \rightarrow \mu x^+$ and $x^- \rightarrow \frac{x^-}{\mu}$, we get the metric as

$$ds^2 = -4dx^+dx^- - \mu^2 \left(x^2 + \frac{R_A^2}{R_S^2}y^2\right)(dx^+)^2 + dx^2 + x^2d\Omega_{p-2}^2 + dy^2 + y^2d\Omega_{q-2}^2. \quad (6)$$

It's easy to see from the Penrose limit of $AdS_p \times S^q$ that the Penrose limit of spaces with different values of p and q but with same p+q have the same pp wave provided the radius of curvature of AdS is equal to the radius of sphere. For $AdS_5 \times S^5$ the corresponding R-R 5-form field strength is

$$F_{+1234} = F_{+5678} = \text{constant} \times \mu. \quad (7)$$

The world volume theory of Dp brane supports solitonic configurations of lower dimensional branes. In flat space, in particular, in the context of D3 branes, BPS monopoles correspond to an orthogonal D1 brane. We shall describe D3 brane in the Penrose limit of $AdS_5 \times S^5$ background.

The low energy dynamics of a D3 brane in a pp wave background, i.e

$$ds^2 = -4dx^+dx^- - \mu^2(x_1^2 + \dots + x_8^2)(dx^+)^2 + \sum_{i,j=1}^8 dx^i dx^j \eta_{ij}, \quad (8)$$

where we have taken the radius of curvature of AdS is same as the radius of the sphere, is described by

$$S = -T_3 \int d^4\sigma \sqrt{|det(P[G]_{ab} + \lambda F_{ab})|}, \quad (9)$$

where we have set $B_{\mu\nu} = 0$, dilaton=0. F_{ab} is the U(1) field strength living on the brane, $\lambda \equiv 2\pi\alpha'$, but we shall set $\lambda = 1$, henceforth, and P is the pullback which will pullback the bulk fields onto the worldvolume of the brane and $P[G]_{ab}$ is given by

$$P[G]_{ab} = -2\partial_a X^+ \partial_b X^- - 2\partial_a X^- \partial_b X^+ - \mu^2(x_1^2 + \dots + x_8^2) \partial_a X^+ \partial_b X^+ + \sum_{i,j=1}^8 \partial_a X^i \partial_b X^j \eta_{ij}. \quad (10)$$

We shall choose the static gauge choice as²

$$X^+ = \phi^9, \quad X^- = \phi^{10}, \quad X^1 = \sigma^1, \quad X^2 = \sigma^2, \quad X^3 = \sigma^3, \quad X^4 = \sigma^4, \quad X^m = \phi^m. \quad (11)$$

²In our notation a,b denotes the worldvolume coordinates of D3 brane and can take values 1,2,3,4 and m can take values from 5 to 8.

In this choice of static gauge pullback becomes

$$P[G]_{ab} = +\partial_a \sigma^1 \partial_b \sigma^1 + \partial_a \sigma^2 \partial_b \sigma^2 + \partial_a \sigma^3 \partial_b \sigma^3 + \partial_a \sigma^4 \partial_b \sigma^4 + \partial_a \phi \partial_b \phi, \quad (12)$$

where we have excited only one transverse scalar, say $\phi = \phi^5$. Let us also excite a magnetic field $B^\alpha = \frac{1}{2}\epsilon^{\alpha\beta\gamma}F_{\beta\gamma}(\alpha, \beta, \gamma = 2, 3, 4)$, and shall take the 'time' as σ^1 . For static configurations, the energy then becomes

$$\begin{aligned} E = -L &= T_3 \int d^3\sigma \left[(1 \pm \vec{B} \cdot \vec{\nabla}\phi)^2 + (\vec{B} \mp \vec{\nabla}\phi)^2 \right]^{\frac{1}{2}} \\ T_3 &\geq \int d^3\sigma (1 \pm \vec{B} \cdot \vec{\nabla}\phi) \end{aligned} \quad (13)$$

The lower bound is achieved when

$$\vec{B} = \pm \vec{\nabla}\phi, \quad (14)$$

using this along with the Bianchi identity, $\vec{\nabla} \cdot \vec{B} = 0$, we get the equation to scalar as

$$\nabla^2 \phi = 0. \quad (15)$$

The trivial solutions to eq.(15) and magnetic field is

$$\phi = \text{constant}, \quad \vec{B} = 0, \quad (16)$$

and the nontrivial solution is

$$\phi = \frac{N}{2r}, \quad \vec{B} = \mp \frac{N}{2r^3} \vec{r}, \quad (17)$$

where $r^2 = (\sigma^1)^2 + (\sigma^2)^2 + (\sigma^3)^2$ and N is an integer due to charge quantization [45]. Hence, the energy to this configuration is the sum of energy of Born-Infeld and Chern-Simon part. Since, we are taking minimum energy for the nonlinear Born-Infeld action implies the total energy is minimum.

In order to see how much supersymmetry does this configuration preserve let us note that the supersymmetric analysis to D3 brane have been studied in [46],[47],[43] and in [40], these authors have noted that D3 brane preserves half of the supersymmetry in the background of pp waves provided the worldvolume directions of D3 are along the light cone directions along with two other directions are along either of SO(4) directions. If one of the direction is along one SO(4) and other is along the other SO(4) direction then the solution do not preserve any supersymmetry [42].

It has been argued that with every brane embedding there is a kappa symmetry projection operator which satisfies

$$\Gamma \epsilon = \epsilon, \quad (18)$$

if the given brane embedding preserves some fraction of the supersymmetry, where ϵ is a Killing spinor of the supersymmetric background and Γ is kappa symmetric projector and is defined as in [48]-[50]

$$d^{p+1}\sigma\Gamma = -\frac{e^{\frac{p-3}{4}\phi}}{\mathcal{L}_{\mathcal{DBI}}} \exp(e^{-\frac{\phi}{2}}\mathcal{F}) \wedge XY|_{vol}, \quad (19)$$

with

$$X = \bigoplus_n \gamma_{(2n+q)} P^{n+q}, \quad (20)$$

where

$$\begin{aligned} IIA : P &= \gamma_{11} \quad Y = 1, \quad q = 1 \\ IIB : P &= K, \quad Y = I, \quad q = 0, \end{aligned} \quad (21)$$

where $|_{vol}$ means that term should be proportional to the volume form of the brane and $\mathcal{F} = F - B$. The operators I and K act on spinors as $I\psi = -i\psi$ and $K\psi = \psi^*$ respectively. These operators are anticommuting and satisfy the following properties: $I^2 = -1$, $J^2 = 1 = K^2$ and $IJ=K$. $\frac{1}{\mathcal{L}_{\mathcal{DBI}}}$ is the value of the DBI Lagrangian evaluated in the background with the embeddings. $\gamma_{(2n)}$ is a 2n-form gamma matrices defined on target space as

$$\gamma_{(2n)} = \frac{1}{2n!} \gamma_{i_1 \dots i_{2n}} d\sigma^{i_1} \wedge \dots \wedge d\sigma^{i_{2n}}, \quad (22)$$

and $\gamma_{i_1 \dots i_{2n}}$ is the pullback of the target space gamma matrices

$$\gamma_{i_1 \dots i_{2n}} = \partial_{i_1} X^{\mu_1} \dots \partial_{i_{2n}} X^{\mu_{2n}} \gamma_{\mu_1 \dots \mu_{2n}}. \quad (23)$$

It has been shown in [48] that Γ satisfies the following properties. $\Gamma^2 = 1$ and $\text{tr}(\Gamma) = 0$, which enables to define projector of that kind which projects out half of the worldvolume fermions thus making the degrees of freedom of worldvolume fermions and bosons same.

Since, we are studying a D3 brane configuration where dilaton and 2-form anti-symmetric B field is zero in a IIB background then the projector written in eq. (19) becomes

$$\Gamma = -\frac{\epsilon^{i_1 \dots i_4}}{\mathcal{L}_{\mathcal{DBI}}} \left[\frac{1}{24} \gamma_{i_1 \dots i_4} I + \frac{1}{4} F_{i_1 i_2} \gamma_{i_3 i_4} J + \frac{1}{8} F_{i_1 i_2} F_{i_3 i_4} I \right]. \quad (24)$$

Evaluating the projector in our static gauge choice, we get:

$$\begin{aligned} \Gamma = & -\frac{i}{\mathcal{L}_{\mathcal{DBI}}} \left[(\gamma_{1234} + \partial_2 \phi \gamma_{1345} - \partial_3 \phi \gamma_{1245} + \partial_4 \phi \gamma_{1235}) I \right. \\ & \left. + \{ B^2 (\gamma_{12} + \partial_2 \phi \gamma_{15}) + B^3 (\gamma_{13} + \partial_3 \phi \gamma_{15}) + B^4 (\gamma_{14} + \partial_4 \phi \gamma_{15}) \} J \right]. \end{aligned} \quad (25)$$

The first and second line of eq. (25) follows from the first and second term of eq. (24) and the last term vanishes because of our choice of magnetic field. For negative chirality type IIB spinors

$$\gamma_{+-12345678} \epsilon = -\epsilon, \quad (26)$$

the Killing spinor, ϵ , is derived by setting the supersymmetric variation of dilatino and gravitino to zero, of the pp-wave is given by [1], [42]

$$\begin{aligned} \epsilon = & \left\{ 1 - \frac{i}{2} \sum_{a=1}^4 \gamma_{-} (y^a \gamma_a \gamma_{1234} + z^a \gamma_{(a+4)} \gamma_{5678}) \right\} \\ & \left(\cos \frac{x^+}{2} - i \sin \frac{x^+}{2} \gamma_{1234} \right) \left(\cos \frac{x^+}{2} - i \sin \frac{x^+}{2} \gamma_{5678} \right) (\lambda + i\eta), \end{aligned} \quad (27)$$

where λ and η are constant, real negative chiral spinors and y^a , and z^a are the coordinates along the first and second $SO(4)$ respectively. Substituting the value of Γ and ϵ in eq. (18), and restricting to $\phi = \phi(\sigma^2)$ and only one component of magnetic field i.e. to B^2 , for simplicity, we get

$$\epsilon = -\frac{1}{\mathcal{L}_{DBI}} \left[(\gamma_{1234} + \partial_2 \phi \gamma_{1345}) \epsilon - B^2 (\gamma_{12} + \partial_2 \phi \gamma_{15}) \epsilon^* \right]. \quad (28)$$

Let us define

$$\begin{aligned} P & \equiv -\frac{1}{2} \sum_{a=1}^4 \gamma_{-} (y^a \gamma_a \gamma_{1234} + z^a \gamma_{(a+4)} \gamma_{5678}) \\ Q & \equiv B^2 (\gamma_{12} + \partial_2 \phi \gamma_{15}) \\ R & \equiv \gamma_{1234} + \partial_2 \phi \gamma_{1345}. \end{aligned} \quad (29)$$

using these, we can rewrite eq. (28) as

$$\epsilon = -\frac{1}{\mathcal{L}_{DBI}} \left[R \epsilon - Q \epsilon^* \right]. \quad (30)$$

Since the Killing spinor eq. (27) holds for all values of x^+ , implies, on plugging into eq. (30) gives rise to

$$\begin{aligned} \left(1 + \frac{R}{\mathcal{L}_{DBI}} \right) (1 + iP) (\lambda + i\eta) &= \frac{Q}{\mathcal{L}_{DBI}} (1 - iP) (\lambda - i\eta) \\ \left(1 + \frac{R}{\mathcal{L}_{DBI}} \right) (1 + iP) (\gamma_{1234} + \gamma_{5678}) (\lambda + i\eta) &= -\frac{Q}{\mathcal{L}_{DBI}} (1 - iP) (\gamma_{1234} + \gamma_{5678}) (\lambda - i\eta) \\ \left(1 + \frac{R}{\mathcal{L}_{DBI}} \right) (1 + iP) \gamma_{1\dots 8} (\lambda + i\eta) &= \frac{Q}{\mathcal{L}_{DBI}} (1 - iP) \gamma_{1\dots 8} (\lambda - i\eta). \end{aligned} \quad (31)$$

Equating real and imaginary parts of eq. (31) and using negative chirality of λ and η , we get

$$\begin{aligned} \left(1 + \frac{R - Q}{\mathcal{L}_{DBI}} \right) (\lambda - P\eta) &= 0, \\ \left(1 + \frac{R + Q}{\mathcal{L}_{DBI}} \right) (P\lambda + \eta) &= 0, \end{aligned}$$

$$\begin{aligned}
\gamma_- \gamma_+ \left(1 + \frac{R-Q}{\mathcal{L}_{\mathcal{DBI}}}\right) \lambda &= 0, \\
\gamma_- \gamma_+ \left(1 + \frac{R+Q}{\mathcal{L}_{\mathcal{DBI}}}\right) \eta &= 0, \\
\gamma_- \gamma_+ \left(1 + \frac{R+Q}{\mathcal{L}_{\mathcal{DBI}}}\right) \gamma_{1234} \lambda &= 0, \\
\gamma_- \gamma_+ \left(1 + \frac{R-Q}{\mathcal{L}_{\mathcal{DBI}}}\right) \gamma_{1234} \eta &= 0.
\end{aligned} \tag{32}$$

Let us see how much supersymmetry does this configuration preserves for zero magnetic field i.e. $B^2 = 0$. From third and fourth equation of eq. (32) we get

$$\begin{aligned}
(1 + \gamma_{1234}) \lambda &= 0, \\
(1 + \gamma_{1234}) \eta &= 0,
\end{aligned} \tag{33}$$

which also follows from fifth and sixth equation of eq. (32). However, from first and second equation of eq. (32) we get on invoking eq. (33)

$$\begin{aligned}
(1 - \gamma_{1234}) \lambda &= 0 \\
(1 - \gamma_{1234}) \eta &= 0.
\end{aligned} \tag{34}$$

Compatibility of eq. (33) and eq. (34) implies that for zero magnetic field the configuration breaks all supersymmetry. Doing the same kind of analysis for nonzero magnetic field we conclude that the configuration breaks all supersymmetry.

Let us look at a different configuration, $(+, 3, 0)$, which means that the worldvolume directions of the brane are extended along the positive lightcone and along one of the $SO(4)$ directions. In this case we shall choose our static gauge choice as

$$X^+ = \tau, \quad X^- = \phi^9, \quad X^1 = \sigma^1, \quad X^2 = \sigma^2, \quad X^3 = \sigma^3, \quad X^4 = \phi^4, \quad X^m = \phi^m. \tag{35}$$

In this choice of static gauge pullback becomes³

$$\begin{aligned}
P[G]_{ab} &= -\mu^2 \left((\sigma^1)^2 + (\sigma^2)^2 + (\sigma^3)^2 + \phi^4 \phi^4 + \phi^m \phi^m \right) \partial_a \tau \partial_b \tau + \partial_a \sigma^1 \partial_b \sigma^1 + \\
&\quad \partial_a \sigma^2 \partial_b \sigma^2 + \partial_a \sigma^3 \partial_b \sigma^3 + \partial_a \phi^4 \partial_b \phi^4 + \partial_a \phi^m \partial_b \phi^m,
\end{aligned} \tag{36}$$

where we have taken the spacetime coordinates as same as the embeddings, i.e. $x^i = X^i$, $i = 1, \dots, 8$. Let us excite only one transverse scalar, say $\phi = \phi^4$ and a magnetic field $B^\alpha = \frac{1}{2} \epsilon^{\alpha\beta\gamma} F_{\beta\gamma}(\alpha, \beta, \gamma = 1, 2, 3)$. For static configurations, the energy then becomes

$$\begin{aligned}
E = -L &= T_3 \int d^3 \sigma \sqrt{\mu^2 ((\sigma^1)^2 + (\sigma^2)^2 + (\sigma^3)^2 + (\phi)^2)} \left[(1 \pm \vec{B} \cdot \vec{\nabla} \phi)^2 + (\vec{B} \mp \vec{\nabla} \phi)^2 \right]^{\frac{1}{2}} \\
T_3 &\geq \int d^3 \sigma \sqrt{\mu^2 ((\sigma^1)^2 + (\sigma^2)^2 + (\sigma^3)^2 + (\phi)^2)} (1 \pm \vec{B} \cdot \vec{\nabla} \phi)
\end{aligned} \tag{37}$$

³Here a,b can take values 0,1,2,3

The lower bound to energy and the solution are written in eq. (14) and eq. (17). However, the consistency of the static gauge choice implies that we have to find solution to ϕ and B by taking the following restriction into account:

$$\mu^2 \left((\sigma^1)^2 + (\sigma^2)^2 + (\sigma^3)^2 + (\phi)^2 \right) = \text{constant} \equiv c^2. \quad (38)$$

So, the solution to the restriction and the equation satisfied by B and ϕ gives us

$$\phi = \text{constant}, \quad B = 0, \quad (\sigma^1)^2 + (\sigma^2)^2 + (\sigma^3)^2 = \text{constant} \equiv c_1^2, \quad (39)$$

by looking at the solution we conclude that the brane is a spherical D3 brane. To check the supersymmetry preserved by this configuration, let us take the projector as

$$\Gamma = -\frac{1}{c}(\gamma_+ - \frac{c^2}{2}\gamma_-)\gamma_{123}I. \quad (40)$$

Substituting the above form of Γ and ϵ from eq. (27) in eq. (18), we get the following equations.

$$\begin{aligned} (1 - i(Q - R))(1 + iP)(\lambda + i\eta) &= 0 \\ (1 - i(Q - R))(1 + iP)\gamma_{1\dots 8}(\lambda + i\eta) &= 0 \\ (1 - i(Q - R))(1 + iP)(\gamma_{1234} + \gamma_{5678})(\lambda + i\eta) &= 0, \end{aligned} \quad (41)$$

where $Q = \frac{\gamma_{+123}}{c}$ and $R = \frac{c}{2}\gamma_{-123}$. Using $\gamma_-^2 = 0$, we get from second and third equation of eq. (41)

$$\gamma_+\lambda = 0, \quad \gamma_+\eta = 0. \quad (42)$$

On invoking this condition on λ and η in the first equation of eq. (41), we see that the configuration preserves one half supersymmetry for $P = -\frac{c}{2}\gamma_{-123}$. However, for $P = -\frac{c}{2}\gamma_{-234}$, we get one more condition on λ and η . Hence, for the later choice of P we conclude that the configuration preserves one quarter supersymmetry. So, finally, for different choices of P the configuration preserves different amount of supersymmetry.

We have derived nonsupersymmetric and supersymmetric solutions of D3 brane in the pp wave background for two different kind of configurations, in one case the brane is extended completely along one of the $SO(4)$ directions and in the other case one of the worldvolume direction is extended along the positive lightcone direction and the rest three are along one of the $SO(4)$ directions. In the former case we found that our solution breaks all supersymmetry for zero magnetic field on the worldvolume of the brane and for a constant transverse scalar and also for nonzero magnetic field and nonconstant transverse scalar. Where as in the later configuration it preserves different amount of supersymmetry depending on the choices of P .

Acknowledgements

I would like to thank Alok kumar for many useful discussions, Sunil Mukhi for many useful correspondences and to the anonymous referee for several constructive suggestions.

References

1. J. Figueroa-O'Farrill, G. Papadopoulos, "Homogeneous Fluxes, Branes and a Maximally Supersymmetric Solution of M-theory," *JHEP* **0108** (2001) 036; hep-th/0105308. M. Blau, J. Figueroa-O'Farrill, C. Hull, G. Papadopoulos, "Penrose Limits and Maximal Supersymmetry," hep-th/0210081. M. Blau, J. Figueroa-O'Farrill, C. Hull, G. Papadopoulos, "A New Maximally Supersymmetric Background of IIB Superstring Theory," *JHEP* **0201** (2002) 047, hep-th/0110242; M. Blau, J. Figueroa-O'Farrill, G. Papadopoulos, "Penrose limits, supergravity and brane dynamics," hep-th/0202111.
2. D. Berenstein, J. Maldacena and H. Nastase, "Strings in flat space and pp waves from $\mathcal{N} = 4$ super Yang Mills," hep-th/0202021.
3. D. Berenstein, E. Gava, J. Maldacena, K.S. Narain and H. Nastase, "Open strings on plane waves and their Yang-Mills duals," hep-th/0203249.
4. C. Kristjansen, J. Plefka, G.W. Semenoff and M. Staudacher, "A New double-scaling limit of N=4 super Yang-Mills theory and pp-wave strings," hep-th/0205033.
5. S.S. Gubser, I.R. Klebanov and A.M. Polyakov, "A semi-classical limit of the gauge/string correspondence," hep-th/0204051.
6. D. Berenstein and H. Nastase, "On lightcone string field theory from super Yang-Mills and holography," hep-th/0205048.
7. D.J. Gross, A. Mikhailov and R. Roiban, "Operators with large R charge in N=4 Yang-Mills theory," hep-th/0205066.
8. N.R. Constable, D.Z. Freedman, M. Headrick, S. Minwalla, L. Motl, A. Postnikov and W. Skiba, "PP-wave string interactions from perturbative Yang-Mills theory," hep-th/0205089.
9. M. Spradlin and A. Volovich, "Superstring interactions in a pp-wave background," hep-th/0204146.
10. R. Gopakumar, "String interactions in pp-waves," hep-th/0205174.
11. M. Cvetič, H. Lu and C.N. Pope, "Penrose limits, pp-waves and deformed M2-branes," hep-th/0203082; "M-theory PP-waves, Penrose Limits and Supernumerary Supersymmetries," hep-th/0203229.

12. H. Singh, “M5-branes with 3/8 supersymmetry in pp wave background,” hep-th/0205020.
13. U. Gursoy, C. Nunez and M. Schvellinger, “RG flows from Spin(7), CY 4-fold and HK manifolds to AdS, Penrose limits and pp waves,” hep-th/0203124.
14. J. Michelson, “(Twisted) toroidal compactification of pp-waves,” hep-th/0203140.
15. M. Hatsuda, K. Kamimura and M. Sakaguchi, “From super- $AdS_5 \times S^5$ algebra to super-pp-wave algebra,” hep-th/0202190; “Super-pp-wave algebra from super- $AdS \times S$ algebras in eleven-dimensions,” hep-th/0204002.
16. J.P. Gauntlett and C.M. Hull, “PP-waves in 11-dimensions with extra supersymmetry,” hep-th/0203255.
17. P. Lee and J. Park, “Open strings in pp-wave background from defect conformal field theory,” hep-th/0203257.
18. V. Balasubramanian, M. Huang, T.S. Levi and A. Naqvi, “Open strings from N=4 super Yang-Mills,” hep-th/0204196.
19. C. Ahn, “More on Penrose limit of $AdS_4 \times Q^{1,1,1}$,” hep-th/0205008; “Comments on Penrose Limit of $AdS_4 \times M^{1,1,1}$,” hep-th/0205109.
20. A. Parnachev and D.A. Sahakyan, “Penrose limit and string quantization in $AdS_3 \times S^3$,” hep-th/0205015.
21. M. Li, “Correspondence principle in a pp-wave background,” hep-th/0205043.
22. S.D. Mathur, A. Saxena and Y.K. Srivastava, “Scalar propagator in the pp-wave geometry obtained from $AdS_5 \times S^5$,” hep-th/0205136.
23. K. Dasgupta, M.M. Sheikh-Jabbari and M. van Raamsdonk, “Matrix perturbation theory for M-theory on a pp-wave,” hep-th/0205185.
24. G. Bonelli, “Matrix strings in pp-wave backgrounds from deformed super Yang-Mills Theory,” hep-th/0205213.
25. Y. Kiem, Y. Kim, S. Lee, J. Park, “PP-wave/Yang-Mills Correspondence: An Explicit Check,” hep-th/0205279.
26. N. Itzhaki, I. R. Klebanov and S. Mukhi, “PP Wave Limit and Enhanced Supersymmetry in Gauge Theories,” hep-th/0202153; S. Mukhi, M. Rangamani and E. Verlinde, “Strings from Quivers, Membranes from Moose,” hep-th/0204174.
27. J. Gomis and H. Ooguri, “Penrose Limit of $\mathcal{N}=1$ Gauge Theories,” hep-th/0202157.

28. L. A. Pando-Zayas and J. Sonnenschein, “On Penrose limits and gauge theories,” hep-th/0202186.
29. M. Alishahia and M. M. Sheikh-Jabbari, “The PP-wave Limit of orbifolded $AdS_5 \times S^5$,” hep-th/0203018; “Strings in PP-waves and Worldsheet Deconstruction,” hep-th/0204174.
30. N. Kim, A. Pankiewicz, S-J. Rey and S. Theisen, “Superstring on PP-Wave Orbifold from Large-N Quiver Theory,” hep-th/0203080.
31. S. R. Das, C. Gomez and S. J. Rey, “Penrose limits, spontaneous symmetry breaking and holography in pp wave background,” hep-th/0203164.
32. R. G. Leigh, K. Okuyama and M. Rozali, “PP-waves and holography,” hep-th/0204026.
33. E. Kiritsis and B. Pioline, “Strings in homogeneous gravitational waves and null holography,” hep-th/0204004.
34. T. Takayanagi and S. Tereshima, “Strings on Orbifolded PP-Waves,” hep-th/0203093.
35. K. Oh and R. Tatar, “Orbifolds, Penrose limits and supersymmetry enhancement,” hep-th/0205067.
36. R. R. Metsaev, “Type IIB Green-Schwarz superstring in plane wave Ramond-Ramond background,” Nucl. Phys. B **625**, 70, (2002) hep-th/0112044.
37. R. R. Metsaev and A. A. Tseytlin, “Exactly solvable model of superstring in Ramond-Ramond background,” hep-th/0202109.
38. J. G. Russo and A. A. Tseytlin, “On solvable models of type IIB superstrings in NS-NS and R-R plane wave backgrounds,” hep-th/0202179.
39. S. Frolov and A.A. Tseytlin, “Semiclassical quantization of rotating superstring in $AdS_5 \times S^5$,” hep-th/0204226.
40. A. Dabholakar and S. Parvizi, “ DP Branes in PP-wave Background,” hep-th/0203231.
41. D. Bak, “Supersymmetric branes in PP Wave Background,” hep-th/0204033.
42. K. Skenderis and M. Taylor, “ Branes in AdS and pp-wave spacetimes,” hep-th/0204054.
43. P. Bain, P. Meessen and M. Zamaklar, “Supergravity solutions for D-branes in Hpp-wave background,” hep-th/0205106.

44. A. Kumar, R. R. Nayak and Sanjay, “D-Brane Solutions in pp-wave Background,” hep-th/0204025; A. Kumar and M. Alishahiha, “D-Brane Solutions from New Isometries of pp-Waves,” hep-th/0205134.
45. N. R. Constable, R. C. Myers and O. Tafjord, “Fuzzy funnels: Non-abelian Brane Intersections,” hep-th/0105035.
46. M. Billo’, I. Pesando, “Boundary States for GS superstrings in an Hpp wave background,” hep-th/0203028.
47. Chong-Sun Chu and Pei-Ming Ho, “Noncommutative D-brane and Open String in pp-wave Background with B-field,” hep-th/0203186.
48. M. Cederwall, A. von Gussich, B. Nilsson and A. Westerberg, “The Dirichlet super-three-brane in ten-dimensional type IIB supergravity,” Nucl. Phys. B **490**, 163 (1997), hep-th/9610148; M. Cederwall, A. von Gussich, P. Sundell and A. Westerberg, “The Dirichlet super-p-branes in ten dimensional type IIA and IIB supergravity,” Nucl. Phys. B **490**, 179, (1997), hep-th/9611159.
49. M. Aganagic, C. Popescu and J. Schwarz, “D-brane actions with local kappa symmetry,” Phys. Lett. B **393**, 311, (1997), hep-th/9610249; “Gauge-invariant and gauge fixed D-brane actions,” Nucl. Phys. B **495**, 99, (1997), hep-th/9612080.
50. E. Bergshoeff and P. Townsend, “Super D-branes,” Nucl. Phys. B **490**, 145, (1997), hep-th/9611173; E. Bergshoeff, R. Kallosh, T. Ortin and G. Papadopoulos, “Kappa-symmetry, supersymmetry and intersecting branes,” Nucl. Phys. B **502**, 149, (1997), hep-th/9705040.